

Past, Present, and Future of Mutagens in Cooked Foods

by Takashi Sugimura*

Mutation assay with *Salmonella typhimurium* enabled us to detect various types of mutagens in cooked foods. A series of mutagenic heterocyclic amines have been isolated and identified in broiled fish and meat and in pyrolyzates of amino acids and proteins. Feeding experiments showed these mutagens to be carcinogenic in mice and rats. The mechanism of formation and pathway of metabolic activation of these heterocyclic amines have been elucidated. Their contents in various cooked foods have been determined. The presence of mutagenic nitropyrenes (some of which were confirmed as carcinogens) in grilled chicken was also established. Roasted coffee beans also yield mutagens such as methylglyoxal. The formation of mutagen precursors, including β -carboline derivatives and tyramine which become mutagens with nitrite treatment, was found during food processing. Oncogene activation in animal tumors induced by some of these food mutagens/carcinogens has been confirmed. The role of mutagens/carcinogens in cooked foods in human cancer development has not yet been exactly evaluated. In order to do this, more information on their carcinogenic potency, human intake, metabolism in the human body, and the effects of combined administration with other initiators, promoters and other modifying factors in food is required.

General Background

It has been generally accepted that the incidence of cancers in different organs varies very much from nation to nation. Studies on immigrants also revealed that different cancer incidence in various areas of the world depends on the life style, including dietary habits. Careful epidemiological studies led us to conclude that the dietary factor was one of the most important causes of human cancers, as important as cigarette smoking (1-3).

Diet influences the incidence of cancer in many ways, directly and indirectly. In one field of research, trials to detect known carcinogens in food were made many years ago. These are described in two papers by Kuratsune (4) and by Lijinsky and Shubik (5) on the detection of carcinogenic aromatic hydrocarbons in biscuits and broiled meat. Carcinogenesis experiments are time consuming and require many animals. Therefore it was very difficult to search for new carcinogens other than those already known, such as aromatic hydrocarbons, mycotoxins and dialkylnitrosamines from foods.

Since the overlapping of mutagens and carcinogens was suggested, especially from data obtained by using the Ames test with *Salmonella typhimurium* (6-8), it became possible to predict the presence of carcinogens in food through the detection of mutagenic activity of food. The formation of mutagens detectable by the Ames test upon broiling dried fish and ground meat was reported by us (9) and Commoner et al. (10). These

studies prompted us to identify mutagens formed during cooking. New mutagens have been discovered successively and these chemicals synthesized and subjected to long-term animal tests. Some of them have now been demonstrated to be carcinogenic in mice and rats (11-15). Thus, the introduction of microbial mutation to screen for carcinogens formed by cooking was proven to be practical and realistic.

Mutagens Found in Cooked Foods and Heated Amino Acids and a Protein

Kasai et al. (16,17) were the first to isolate and identify aminoimidazoquinoline (IQ) and aminomethylimidazoquinoline (MeIQ) from broiled sun-dried sardines (Table 1). They (18) also isolated aminomethylimidazoquinoxaline (MeIQx) from fried beef. The contributions of the Maillard reaction and of the presence of creatinine in meat to the formation of aminoimidazoquinolines and aminomethylimidazoquinoxalines, were proposed (19). Actually, the formation of MeIQx and DiMeIQx after heating of the mixtures of creatinine, sugars and amino acids was demonstrated by Jägerstad et al. (20) and Negishi et al. (21,22). The specific mutagenic activities of IQ, MeIQ, and MeIQx are very high towards *S. typhimurium* TA 98 with S-9 mix as shown in Table 2. Ohgaki et al. demonstrated the carcinogenicity of IQ (23) and MeIQ (24), in mice while Takayama et al. (25) and Tanaka et al. (26) proved the carcinogenicity of IQ in rats. The locations of tumors in mice

*National Cancer Center, 1-1, Tsukiji 5-chome, Chuo-ku, Tokyo 104, Japan.




Table 1. Abbreviations used.

Abbreviation	Compound
IQ	2-Amino-3-methylimidazo[4,5-f]quinoline
MeIQ	2-Amino-3,4-dimethylimidazo[4,5-f]quinoline
MeIQx	2-Amino-3,8-dimethylimidazo[4,5-f]quinoxaline
4,8-DiMeIQx	2-Amino-3,4,8-trimethylimidazo[4,5-f]quinoxaline
7,8-DiMeIQx	2-Amino-3,7,8-trimethylimidazo[4,5-f]quinoxaline
Trp-P-1	3-Amino-1,4-dimethyl-5H-pyrido[4,3-b]indole
Trp-P-2	3-Amino-1-methyl-5H-pyrido[4,3-b]indole
Glu-P-1	2-Amino-6-methyldipyrido[1,2-a:3',2'-d]imidazole
Glu-P-2	2-Aminodipyrido[1,2-a:3',2'-d]imidazole
AαC	2-Amino-9H-pyrido[2,3-b]indole or 2-amino-α-carboline
MeAαC	2-Amino-3-methyl-9H-pyrido[2,3-b]indole or 2-amino-3-methyl-α-carboline

Table 2. Mutagenic activities of aminoimidazoquinolines and aminoimidazoquinoxalines toward *S. typhimurium* with S-9 mix.

	Mutagenic activity, revertants/μg	
	TA 98	TA 100
IQ	433,000	7,000
MeIQ	661,000	30,000
MeIQx	145,000	14,000
4,8-DiMeIQx	183,000	8,000
7,8-DiMeIQx	163,000	9,900

Table 3. Location of tumors induced by IQ and MeIQ in mice and rats.

	Mouse	Rat
IQ	Liver Forestomach Lung	Liver Small and large intestines Zymbal gland Clitoral gland Skin Oral cavity
MeIQ	Liver Forestomach Small and large intestines	Not done

and rats are given in Table 3. The metabolic conversion of the amino group of these heterocyclic amines to the hydroxyamino group catalyzed by cytochrome P-450s is essential to the exertion of their mutagenic activities.

Some heterocyclic amines are isolated from pyrolyzates of amino acids and a protein as a model system of charred food. Thus, Trp-P-1 and Trp-P-2 from a tryptophan pyrolyzate were reported by Sugimura et al. (27) and Glu-P-1 and Glu-P-2 from a glutamic acid pyrolyzate were reported by Yamamoto et al. (28). Amino-α-carboline (AαC) and methylamino-α-carboline (MeAαC) were isolated from a pyrolyzate of soybean globulin by Yoshida et al. (29). The presence of Glu-P-2 (30), Trp-P-1 and Trp-P-2 (31), and AαC and MeAαC in broiled foods (32) and of AαC and MeAαC in cigarette smoke condensates (32,33) has been reported. These heterocyclic amines also possess a fairly high mutagenic activity toward *S. typhimurium* TA 98 with S9 mix as shown in Table 3. Trp-P-1 and Trp-P-2 were demonstrated to be carcinogenic in mice by Matsukura et al. (34) and in rats by Hosaka et al. (35) and Takayama et al. (36). Carcinogenicity of Glu-P-1 and Glu-P-2 was described in mice by Ohgaki et al. (37) and in rats by Takayama et al. (38). AαC and MeAαC were carcinogenic in mice (37), but experiments on rats are being repeated due to less carcinogenic potential of AαC and greater toxic potential of MeAαC than shown in mice; in rats MeAαC induces atrophy of the salivary glands and pancreatic acinar cells, as observed by Takayama et al. (39). The locations of tumors in mice and rats fed on diets containing these heterocyclic amines are listed in Table 5.

These heterocyclic amines are metabolized to *N*-hydroxyamino derivatives, and acetyl or sulfate esters would be the ultimate forms which react with DNA as shown by Yamazoe et al. (40,41), Hashimoto et al. (42,43), and Nagao et al. (44).

Tsuda et al. found that Trp-P-1, Trp-P-2, Glu-P-1, Glu-P-2, AαC, and MeAαC were sensitive to nitrite treatment, while IQ, MeIQ, MeIQx, and DiMeIQx are resistant, and that both groups of heterocyclic amines lost their mutagenic activity after treatment with hypochlorite (45-47). Differential inactivation of mutagenic activity by these treatments enabled us to estimate the contributions of IQ-type and non-IQ-type

Table 4. Mutagenic activities of heterocyclic amines isolated from pyrolyzates of amino acids and a protein toward *S. typhimurium* with S-9 mix.

	Mutagenic activity, revertants/ μ g	
	TA 98	TA 100
Trp-P-1	39,000	1,700
Trp-P-2	104,200	1,800
Glu-P-1	49,000	3,200
Glu-P-2	1,900	1,200
AaC	300	20
MeAaC	200	120

heterocyclic amines to the total mutagenicity in basic fractions from various materials. For instance, the mutagenic activity of charred parts of broiled beef consists mainly of IQ-type heterocyclic amines but that of cigarette smoke condensate consists mainly of non-IQ-type mutagens (15).

Quantitative determinations of heterocyclic amines in cooked foods are reported by Felton et al. (48), Hargraves and Pariza (49), Hayatsu et al. (50), and Turesky et al. (51). Recently Grivas and Nyhammar (52) and Takahashi et al. (53) developed a new method for quantitative determination of heterocyclic amines by HPLC with electrochemical detection. This method increased the sensitivity and, with appropriate prepurification steps, should provide us with a more precise value for amounts of heterocyclic amines in various materials. The recoveries of heterocyclic amines during the procedure were determined by using ^{14}C -labeled compounds (53).

Recently Nagao and her associates (54) observed the activation of Ha-ras oncogene in a rat hepatoma which was induced by IQ feeding. They also found an unknown oncogene, other than ras-family or *erbB*, in rat hepatomas induced by IQ. This oncogene can transform NIH 3T3 cells.

Other Mutagens Formed during Cooking

Recently Ohnishi et al. (55) observed the formation of mono- and dinitropyrenes in grilled chicken. They

Table 5. Location of tumors in mice and rats induced by Trp-P-1, Trp-P-2, Glu-P-1, Glu-P-2, AaC, and MeAaC.

	Mouse	Rat
Trp-P-1	Liver	Liver
Trp-P-2	Liver	Liver
Glu-P-1	Liver	Liver
	Blood vessel	Small and large intestines
		Zymbal gland
		Clitoral gland
Glu-P-2	Liver	Liver
	Blood vessel	Small and large intestines
		Zymbal gland
		Clitoral gland
AaC	Liver	Liver
	Blood vessel	
MeAaC	Liver	Not done
	Blood vessel	

calculated that the oral intake of nitropyrenes is much higher than that by inhalation of air polluted with diesel engine exhaust. Nitropyrenes are supermutagens and the carcinogenicity of dinitropyrenes was well documented by Ohgaki et al. (56,57), Tokiwa et al. (58), and Takayama et al. (59).

Nagao and her associates (60) found the activated Ki-ras gene in a rat fibrosarcoma, induced by repeated subcutaneous injections of 1,8-dinitropyrene.

A wide variety of mutagens are produced as shown by early observations made by Kuratsune (4) and Lijinsky and Shubik (5). As an example, it could be mentioned that roasting of coffee beans causes mutagenic activity (61,62). Some portion of the mutagenicity can be accounted for by methylglyoxal (63,64). The production of hydrogen peroxide in brewed coffee and instant coffee solutions was observed under aerobic conditions (65). The co-presence of hydrogen peroxide and methylglyoxal showed enhanced mutagenicity.

Mutagen Precursors Found in Fermented Foods and Vegetables

Piacek-Llanes and Tannebaum (66) first mentioned the presence of a nitrosatable mutagen precursor in fava beans. They postulated a possible relationship between the high incidence of gastric cancers and the intake of fava beans and nitrite in Central and South American countries and searched for mutagens produced by nitrite treatment. The mutagen precursor was later identified as 4-chloro-6-methoxyindole by Yang et al. (67). Soybeans were treated by nitrite but did not exhibit any mutagenic activity, while fermented products of soybeans showed mutagenic activity after nitrite treatment. Fermentation is not "cooking" in a strict sense but is linked in various ways with preparation of food.

Two isomers of 1-methyl-1,2,3,4-tetrahydro- β -carboline-3-carboxylic acids which can be produced by condensation of L-tryptophan and acetaldehyde were found to be mutagen precursors in soy sauce by Wakabayashi et al. (68). Tyramine is also found in soy sauce and a mutagenic diazo-oxide compound was produced after nitrite treatment (69). A similar diazonium compound was reported to be formed from agalitin in mushrooms and reported to produce tumors in the glandular stomach of mice by Toth et al. (70).

From Chinese cabbages, a mutagen precursor was isolated and identified as indole-3-acetonitrile (71,72). Pickling procedures may convert nitrate to nitrite and yield acidic conditions, under which the conversion of a mutagen precursor to a mutagen may occur.

Measures To Be Taken for Cancer Prevention with Regard to Mutagen/Carcinogen Formation during Cooking

Mutagens formed during cooking have been newly discovered by adopting the microbial assay. Their struc-

tures were established and they have been synthesized in quantity by organic chemistry. Long-term carcinogenesis experiments gave positive results for compounds tested so far. An important question arises, namely; how significant are these mutagens in human cancer development?

We need to consider that the carcinogenic process is not a simple mutagenic event. In the case of human carcinogenesis, cancers develop over a long latent period after exposure to mutagenic or carcinogenic factors. An obvious example is that atomic bomb survivors developed leukemia, thyroid cancers, and other cancers many years later. Some subjects who accidentally inhaled mustard gas developed bronchial cancer over 10 years later. In experimental systems also, the process of carcinogenesis is composed of multiple steps. Initiation, which may involve a mutational event, and promotion, which follows the initiation step, lead to the completion of carcinogenesis as proven on the mouse skin by Berenblum (73). The promotion step on mouse skin is catalyzed by tumor promoters including tetradecanoyl phorbol acetate (TPA) (74), teleocidin (75), and aplysiatoxin (76). The multiple steps of promotion and further progression are proposed for the completion of the carcinogenic processes (77). The effects of initiation on various target organs could be readily multiplied and intensified by various factors. Therefore the data on the carcinogenic potential obtained from long-term experiments with rodents given a diet containing a single mutagen can hardly be related directly to human cases. In addition, humans might be different from rodents in their metabolism, repair of DNA modification induced by mutagens and overall susceptibility to carcinogens. Because the completion of the carcinogenic process is the appearance of a tumor arising monoclonally from a single cell, the size of target organs, namely the number of cells existing in target organs should be considered as an important element for more precise risk estimation.

Taking various factors into consideration, it is probably impractical and not realistic to make risk estimations from the carcinogenicity data on rodents given a single carcinogen. However, for a simple extrapolation of animal data for risk estimation, TD_{50} values, which are the doses needed to develop cancers in 50% of animals fed on carcinogens for their life time, have been calculated based on mouse experiments. Those for IQ, Trp-P-1, Trp-P-2, Glu-P-1, Glu-P-2, A α C, and MeA α C were 14.7, 8.8, 2.7, 2.7, 4.9, 15.8, and 5.8 mg/kg/day, respectively. If we assume that the average TD_{50} value of heterocyclic amines should be around 8 mg/kg/day, we can roughly estimate the risk of these carcinogenic heterocyclic amines for human beings. The intake of heterocyclic amines was calculated from available data on their quantities in foods. Apparently the human intake is about 0.0002 times the TD_{50} obtained from animal data. This means that heterocyclic amines may not be so serious for human cancer development. On the other hand, it is also true that humans are being exposed to many heterocyclic amines and many other carcino-

gens with tumor promoters and/or suppressing factors for carcinogenesis. At this moment, it is honest to state that no solid information on the estimation of risk of heterocyclic amines has been obtained in any direction, either positive or negative. Recently Takayama et al. (unpublished data) found that rats given, simultaneously, five heterocyclic amines—IQ, Trp-P-1, Trp-P-2, Glu-P-2, and A α C—each at a dose of one-fifth those given in independent animal experiments resulted in the production of more multiple colon cancers in a shorter period. It suggests to us that simultaneous exposure to different kinds of carcinogens may yield different carcinogenic responses from those of administrations of single carcinogens. We need further experiments using simultaneous administration of different kinds of carcinogens and those using both carcinogen and tumor promoter treatments.

It should be mentioned that the mutation assay systems commonly adopted are able to pick up carcinogens with relatively high mutagenic activity but miss those with low or no mutagenicity. Mutagenicity cannot be a criterion for the detection of tumor promoters which are probably present in food, and for which other methods should be adopted.

Finally, it would be appropriate to establish a way to avoid the formation of mutagens and carcinogens during cooking by improving heating conditions or by including substance(s) suppressing mutagen formation during cooking.

Our work cited in this article was supported by Grants-in-Aid for Cancer Research from the Ministry of Health and Welfare, the Ministry of Education, Science and Culture, Japan, a Grant-in-Aid from the Ministry of Health and Welfare for Comprehensive 10-Year Strategy for Cancer Control, Japan and a grant from Toyota Foundation.

REFERENCES

- Wynder, E. L., and Gori, G. B. Contribution of the environment to cancer incidence: an epidemiologic exercise. *J. Natl. Cancer Inst.* 58: 825–832 (1977).
- Higginson, J., and Muir, C. S. Environmental carcinogenesis: misconceptions and limitations to cancer control. *J. Natl. Cancer Inst.* 63: 1291–1298 (1979).
- Doll, R., and Peto, R. The causes of cancer: quantitative estimates of avoidable risks of cancer in the United States today. *J. Natl. Cancer Inst.* 66: 1193–1308 (1981).
- Kuratsune, M. Benzo[a]pyrene content of certain pyrogenic materials. *J. Natl. Cancer Inst.* 16: 1485–1496 (1956).
- Lijinsky, W., and Shubik, P. Benzo[a]pyrene and other polynuclear hydrocarbons in charcoal-broiled meat. *Science* 145: 53–54 (1964).
- McCann, J., and Ames, B. N. Detection of carcinogens as mutagens in the *Salmonella*/microsome test: assay of 300 chemicals: discussion. *Proc. Natl. Acad. Sci. (U.S.)* 73: 950–954 (1976).
- Purchase, I. F. H., Longstaff, E., Ashby, J., Styles, J. A., Anderson, D., Lefevre, P. A., and Westwood, F. R. Evaluation of six short-term tests for detecting organic chemical carcinogens and recommendations for their use. *Nature* 264: 624–627 (1976).
- Sugimura, T., Sato, S., Nagao, M., Yahagi, T., Matsushima, T., Seino, Y., Takeuchi, M., and Kawachi, T. Overlapping of carcinogens and mutagens. In: *Fundamentals in Cancer Prevention* (P. N. Magee, S. Takayama, T. Sugimura, and T. Matsushima, Eds.), Japan Scientific Societies Press, Tokyo, 1976, pp. 191–215.
- Sugimura, T., Nagao, M., Kawachi, T., Honda, M., Yahagi, T., Seino, Y., Sato, S., Matsukura, N., Matsushima, T., Shirai, A.,

- Sawamura, M., and Matsumoto, H. Mutagen-carcinogens in food, with special reference to highly mutagenic pyrolytic products in broiled foods. In: *Origins of Human Cancer*, Book C (H. H. Hiatt, J. D. Watson, and J. A. Winsten, Eds.), Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1977, pp. 1561-1576.
10. Commoner, B., Vithayathil, A. J., Dolara, P., Nair, S., Madayastha, P., and Cuca, G. C. Formation of mutagens in beef and beef extract during cooking. *Science* 201: 913-916 (1978).
 11. Sugimura, T. Carcinogenicity of mutagenic heterocyclic amines formed during cooking process. *Mutat. Res.* 150: 33-41 (1985).
 12. Sugimura, T., and Sato, S. Mutagen in foods. In: *Voeding en Kanker* (F. H. Sobels and C. M. A. W. Festen, Eds.), Noord-Hollandsche Uitgevers Maatschappij, Amsterdam, 1984, pp. 5-14.
 13. Sugimura, T. Mutagens, carcinogens, and tumor promoters in our daily food. *Cancer* 49: 1970-1984 (1982).
 14. Sugimura, T., and Sato, S. Mutagen-carcinogen in food. *Cancer Res. (Suppl.)* 43: 2415s-2421s (1983).
 15. Sugimura, T., and Sato, S. Bacterial mutagenicity of natural materials, pyrolysis products and additives in foodstuffs and their association with genotoxic effects in mammals. In: *Developments in the Science and Practice of Toxicology* (A. W. Hayes, R. C. Schnell, and T. S. Miya, Eds.), Elsevier, Amsterdam, 1983, pp. 115-133.
 16. Kasai, H., Yamaizumi, Z., Wakabayashi, K., Nagao, M., Sugimura, T., Yokoyama, S., Miyazawa, T., Spingarn, N. E., Weisburger, J. H., and Nishimura, S. Potent novel mutagens produced by broiling fish under normal conditions. *Proc. Japan. Acad.* 56B: 278-283 (1980).
 17. Kasai, H., Yamaizumi, Z., Wakabayashi, K., Nagao, M., Sugimura, T., Yokoyama, S., Miyazawa, T., and Nishimura, S. Structure and chemical synthesis of Me-IQ, a potent mutagen isolated from broiled fish. *Chem. Letters* 1980: 1391-1394 (1980).
 18. Kasai, H., Yamaizumi, Z., Shiomi, T., Yokoyama, S., Miyazawa, T., Wakabayashi, K., Nagao, M., Sugimura, T., and Nishimura, S. Structure of a potent mutagen isolated from fried beef. *Chem. Letters* 1981: 485-488 (1981).
 19. Jägerstad, M., Laser Reuterswärd, A., Öste, R., and Dahlqvist, A. Creatinine and Maillard reaction products as precursors of mutagenic compounds formed in fried beef. In: *The Maillard Reaction in Foods and Nutrition* (G. R. Waller and M. S. Feather, Eds.) (ACS Symposium Series 215), American Chemical Society, Washington, DC, 1983, pp. 507-519.
 20. Jägerstad, M., Olsson, K., Grivas, S., Negishi, C., Wakabayashi, K., Tsuda, M., Sato, S., and Sugimura, T. Formation of 2-amino-3,8-dimethylimidazo[4,5-f]quinoxaline in a model system by heating creatinine, glycine and glucose. *Mutat. Res.* 126: 239-244 (1984).
 21. Negishi, C., Wakabayashi, K., Tsuda, M., Sato, S., Sugimura, T., Saitô, H., Maeda, M., and Jägerstad, M. Formation of 2-amino-3,7,8-trimethylimidazo[4,5-f]quinoxaline, a new mutagen, by heating a mixture of creatinine, glucose and glycine. *Mutat. Res.* 140: 55-59 (1984).
 22. Negishi, C., Wakabayashi, K., Yamaizumi, J., Saitô, H., Sato, S., Sugimura, T., and Jägerstad, M. Identification of 4,8-Di-MeIQx, a new mutagen. *Mutat. Res.* 147: 267-268 (1985).
 23. Ohgaki, H., Kusama, K., Matsukura, N., Morino, K., Hasegawa, H., Sato, S., Sugimura, T., and Takayama, S. Carcinogenicity in mice of a mutagenic compound, 2-amino-3-methylimidazo[4,5-f]quinoline, from broiled sardine, cooked beef and beef extract. *Carcinogenesis* 5: 921-924 (1984).
 24. Ohgaki, H., Hasegawa, H., Kato, T., Suenaga, M., Ubukata, M., Sato, S., Takayama, S., and Sugimura, T. Induction of tumors in the forestomach and liver of mice by feeding 2-amino-3,4-dimethylimidazo[4,5-f]quinoline (MeIQ). *Proc. Japan. Acad.* 61B: 137-139 (1985).
 25. Takayama, S., Nakatsuru, Y., Masuda, M., Ohgaki, H., Sato, S., and Sugimura, T. Demonstration of carcinogenicity in F344 rats of 2-amino-3-methylimidazo[4,5-f]quinoline from broiled sardine, fried beef and beef extract. *Gann* 75: 467-470 (1984).
 26. Tanaka, T., Barnes, W. S., Williams, G. M., and Weisburger, J. H. Multipotential carcinogenicity of the fried food mutagen 2-amino-3-methylimidazo[4,5-f]quinoline in rats. *Japan. J. Cancer Res. (Gann)* 76: 570-576 (1985).
 27. Sugimura, T., Kawachi, T., Nagao, M., Yahagi, T., Seino, Y., Okamoto, T., Shudo, K., Kosuge, T., Tsuji, K., Wakabayashi, K., Iitaka, Y., and Itai, A. Mutagenic principle(s) in tryptophan and phenylalanine pyrolysis products. *Proc. Japan. Acad.* 53B: 58-61 (1977).
 28. Yamamoto, T., Tsuji, K., Kosuge, T., Okamoto, T., Shudo, K., Takeda, K., Iitaka, Y., Yamaguchi, K., Seino, Y., Yahagi, T., Nagao, M., and Sugimura, T. Isolation and structure determination of mutagenic substances in L-glutamic acid pyrolysate. *Proc. Japan. Acad.* 54B: 248-250 (1978).
 29. Yoshida, D., Matsumoto, T., Yoshimura, R., and Matsuzaki, T. Mutagenicity of amino- α -carbolines in pyrolysis products of soybean globulin. *Biochem. Biophys. Res. Commun.* 83: 915-920 (1978).
 30. Yamaguchi, K., Shudo, K., Okamoto, T., Sugimura, T., and Kosuge, T. Presence of 2-aminodipyrido[1,2-a:3',2'-d]imidazole in broiled cuttlefish. *Gann* 71: 743-744 (1980).
 31. Yamaizumi, Z., Shiomi, T., Kasai, H., Nishimura, S., Takahashi, Y., Nagao, M., Sugimura, T. Detection of potent mutagens, Trp-P-1 and Trp-P-2, in broiled fish. *Cancer Letters* 9: 75-83 (1980).
 32. Matsumoto, T., Yoshida, D., and Tomita, H. Determination of mutagens, amino- α -carbolines in grilled foods and cigarette smoke condensate. *Cancer Letters* 12: 105-110 (1981).
 33. Yoshida, D., and Matsumoto, T. Amino- α -carbolines as mutagenic agents in cigarette smoke condensate. *Cancer Letters* 10: 141-149 (1980).
 34. Matsukura, N., Kawachi, T., Morino, K., Ohgaki, H., Sugimura, T., and Takayama, S. Carcinogenicity in mice of mutagenic compounds from a tryptophan pyrolyzate. *Science* 213: 346-347 (1981).
 35. Hosaka, S., Matsushima, T., Hirono, I., and Sugimura, T. Carcinogenic activity of 3-amino-1-methyl-5H-pyrido-[4,3-b]indole(Trp-P-2), a pyrolysis product of tryptophan. *Cancer Letters* 13: 23-28 (1981).
 36. Takayama, S., Nakatsuru, Y., Ohgaki, H., Sato, S., and Sugimura, T. Carcinogenicity in rats of a mutagenic compound, 3-amino-1,4-dimethyl-5H-pyrido[4,3-b]indole, from tryptophan pyrolysate. *Japan. J. Cancer Res. (Gann)*, 76: 815-817 (1985).
 37. Ohgaki, H., Matsukura, N., Morino, K., Kawachi, T., Sugimura, T., and Takayama, S. Carcinogenicity in mice of mutagenic compounds from glutamic acid and soybean globulin pyrolysates. *Carcinogenesis* 5: 815-819 (1984).
 38. Takayama, S., Matsuda, M., Mogami, M., Ohgaki, H., Sato, S., and Sugimura, T. Induction of cancers in the intestine, liver and various other organs of rats by feeding mutagens from glutamic acid pyrolysate. *Gann* 75: 207-213 (1984).
 39. Takayama, S., Nakatsuru, Y., Ohgaki, H., Sato, S., and Sugimura, T. Atrophy of salivary glands and pancreas of rats fed a diet with aminomethyl- α -carboline. *Proc. Japan. Acad.* 61: 277-280 (1985).
 40. Yamazoe, Y., Ishii, K., Kamataki, T., Kato, R., and Sugimura, T. Isolation and characterization of active metabolites of tryptophan pyrolysate mutagen, Trp-P-2, formed by rat liver microsomes. *Chem.-Biol. Interact.* 30: 125-138 (1980).
 41. Yamazoe, Y., Tada, M., Kamataki, T., and Kato, R. Enhancement of binding of N-hydroxy-Trp-P-2 to DNA seryl tRNA synthetase. *Biochem. Biophys. Res. Commun.* 102: 432-439 (1981).
 42. Hashimoto, Y., Shudo, K., and Okamoto, T. Metabolic activation of a mutagen, 2-amino-6-methyldipyrido[1,2-a:3',2'-d]imidazole. Identification of 2-hydroxyamino-6-methyldipyrido[1,2-a:3',2'-d]imidazole and its reaction with DNA. *Biochem. Biophys. Res. Commun.* 92: 971-976 (1980).
 43. Hashimoto, Y., Shudo, K., and Okamoto, T. Activation of a mutagen, 3-amino-1-methyl-5H-pyrido[4,3-b]indole. Identification of 3-hydroxyamino-1-methyl-5H-pyrido[4,3-b]indole and its reaction with DNA. *Biochem. Biophys. Res. Commun.* 96: 355-362 (1980).
 44. Nagao, M., Fujita, Y., Wakabayashi, K., and Sugimura, T. Ultimate forms of mutagenic and carcinogenic heterocyclic amines produced by pyrolysis. *Biochem. Biophys. Res. Commun.* 114: 626-631 (1983).
 45. Tsuda, M., Takahashi, Y., Nagao, M., Hirayama, T., and Sugimura, T. Inactivation of mutagens from pyrolysates of tryptophan

- and glutamic acid by nitrite in acidic solution. *Mutat. Res.* 78: 331-339 (1980).
46. Tsuda, M., Nagao, M., Hirayama, T., and Sugimura, T. Nitrite converts 2-amino- α -carboline, an indirect mutagen, into 2-hydroxy- α -carboline, a non-mutagen, and 2-hydroxy-3-nitroso- α -carboline, a direct mutagen. *Mutat. Res.* 83: 61-68 (1981).
 47. Tsuda, M., Wakabayashi, K., Hirayama, T., and Sugimura, T. Inactivation of potent pyrolysis mutagens by chlorinated tap water. *Mutat. Res.* 119: 27-34 (1983).
 48. Felton, J. S., Knize, M. G., Wood, C., Wuebbles, B. J., Healy, S. K., Stuermer, D. H., Bjeldanes, L. F., Kimble, B. J., and Hatch, F. T. Isolation and characterization of new mutagens from fried ground beef. *Carcinogenesis* 5: 95-102 (1984).
 49. Hargraves, W. A., and Pariza, M. W. Purification and mass spectral characterization of bacterial mutagens from commercial beef extract. *Cancer Res.* 43: 1467-1472 (1983).
 50. Hayatsu, H., Matsui, Y., Ohara, Y., Oka, T., and Hayatsu, T. Characterization of mutagenic fractions in beef extract and in cooked ground beef. Use of blue-cotton for efficient extraction. *Gann* 74: 472-482 (1983).
 51. Turesky, R. J., Wishnok, J. S., Tannenbaum, S. R., Pfund, R. A., and Buchi, G. H. Qualitative and quantitative characterization of mutagens in commercial beef extract. *Carcinogenesis* 4: 863-866 (1983).
 52. Grivas, S., and Nyhammer, T. Analysis of mutagenic imidazo[4,5-f]quinolines and -quinoxalines (IQ compounds). Comparison of electrochemical and ultraviolet detection. *Mutat. Res.* 142: 5-8 (1985).
 53. Takahashi, M., Wakabayashi, K., Nagao, M., Yamamoto, M., Masui, T., Goto, T., Kinae, N., Tomita, I., and Sugimura, T. Quantification of 2-amino-3-methylimidazo[4,5-f]quinoline (IQ) and 2-amino-3,8-dimethylimidazo[4,5-f]quinoxaline (MeIQx) in beef extracts by liquid chromatography with electrochemical detection (LCEC). *Carcinogenesis* 6: 1195-1199 (1985).
 54. Ishikawa, F., Takaku, F., Nagao, M., Ochiai, M., Hayashi, K., Takayama, S., and Sugimura, T. Activated oncogenes in a rat hepatocellular carcinoma induced by 2-amino-3-methylimidazo[4,5-f]quinoline. *Japan. J. Cancer Res. (Gann)* 76: 425-428 (1985).
 55. Ohnishi, Y., Kinouchi, T., Manabe, Y., Tsutsui, H., Otsuka, H., Tokiwa, H., and Otofujii, T. Nitro compounds in environmental mixtures and foods. In: *Short Term Genetics Bioassays in the Evaluation of Complex Environmental Mixtures* (M. D. Waters, S. S. Sandhu, J. Lewtas, L. Claxton, G. Strauss, and S. Nesnow, Eds.), Plenum Press, New York, 1985, pp. 195-204.
 56. Ohgaki, H., Negishi, C., Wakabayashi, K., Kusama, K., Sato, S., and Sugimura, T. Induction of sarcomas in rats by subcutaneous injection of dinitropyrenes. *Carcinogenesis* 5: 583-585 (1984).
 57. Ohgaki, H., Hasegawa, H., Kato, T., Negishi, C., Sato, S., and Sugimura, T. Absence of carcinogenicity of 1-nitropyrene, correction of previous results, and new demonstration of carcinogenicity of 1,6-dinitropyrene in rats. *Cancer Letters* 25: 239-245 (1985).
 58. Tokiwa, H., Otofujii, T., Horikawa, K., Kitamori, S., Otsuka, H., Manabe, Y., Kinouchi, T., and Ohnishi, Y. 1,6-Dinitropyrene: mutagenicity in *Salmonella* and carcinogenicity in BALB/c mice. *J. Natl. Cancer Inst.* 73: 1359-1363 (1984).
 59. Takayama, S., Ishikawa, T., Nakajima, H., and Sato, S. Lung carcinoma induction in Syrian golden hamsters by intratracheal instillation of 1,6-dinitropyrene. *Japan J. Cancer Res. (Gann)* 76: 457-461 (1985).
 60. Ochiai, M., Nagao, M., Tahira, T., Ishikawa, F., Hayashi, K., Ohgaki, H., Tsuchida, N., Terada, M., and Sugimura, T. Activation of K-ras and oncogenes other than ras family, in rat fibro sarcomas initiated by 1,8-dinitropyrene. *Cancer Letters* 29: 119-125 (1985).
 61. Nagao, M., Takahashi, Y., Yamanaka, H., and Sugimura, T. Mutagens in coffee and tea. *Mutat. Res.* 68: 101-106 (1979).
 62. Kosugi, A., Nagao, M., Suwa, Y., Wakabayashi, K., and Sugimura, T. Roasting coffee beans produces compounds that induce prophage λ in *E. coli* and are mutagenic in *E. coli* and *S. typhimurium*. *Mutat. Res.* 116: 179-184 (1983).
 63. Kasai, H., Kumeno, K., Yamaizumi, Z., Nishimura, S., Nagao, M., Fujita, Y., Sugimura, T., Nukaya, H., and Kosuge, T. Mutagenicity of methylglyoxal in coffee. *Gann* 73: 681-683 (1982).
 64. Fujita, Y., Wakabayashi, K., Nagao, M., and Sugimura, T. Characteristics of major mutagenicity of instant coffee. *Mutat. Res. Letters*, in press.
 65. Nagao, M., Suwa, Y., Yoshizumi, H., and Sugimura, T. Mutagens in coffee. *Banbury Report 17: Coffee and Health* (B. MacMahon and T. Sugimura, Eds.), Cold Spring Harbor Lab., Cold Spring Harbor, NY, 1984, pp. 69-77.
 66. Piacek-Llanes, B. G., and Tannenbaum, S. R. Formation of an activated N-nitroso compound in nitrite-treated fava beans (*Vicia faba*). *Carcinogenesis* 3: 1379-1384 (1982).
 67. Yang, D., Tannenbaum, S. R., Büchi, G., and Lee, G. C. M. 4-Chloro-6-methoxyindole is the precursor of a potent mutagen (4-chloro-6-methoxy-2-hydroxy-1-nitroso-indolin-3-one oxime) that forms during nitrosation of the fava beans (*Vicia faba*). *Carcinogenesis* 5: 1219-1224 (1984).
 68. Wakabayashi, K., Ochiai, M., Saitô, H., Tsuda, M., Suwa, Y., Nagao, M., and Sugimura, T. Presence of 1-methyl-1,2,3,4-tetrahydro- β -carboline-3-carboxylic acid, a precursor of a mutagenic nitroso compound, in soy sauce. *Proc. Natl. Acad. Sci. (U.S.)* 80: 2912-2916 (1983).
 69. Ochiai, M., Wakabayashi, K., Nagao, M., and Sugimura, T. Tyramine is a major mutagen precursor in soy sauce, being convertible to a mutagen by nitrite. *Gann* 75: 1-3 (1984).
 70. Toth, B., Nagel, D., and Ross, A. Gastric tumorigenesis by a single dose of 4-(hydroxymethyl)benzenediazonium ion of *Agaricus bisporus*. *Brit. J. Cancer* 46: 417-422 (1982).
 71. Wakabayashi, K., Nagao, M., Ochiai, M., Tahira, T., Yamaizumi, Z., and Sugimura, T. A mutagen precursor in Chinese cabbage, indole-3-acetonitrile, which becomes mutagenic on nitrite treatment. *Mutat. Res. Letters* 143: 17-21 (1985).
 72. Wakabayashi, K., Nagao, M., Tahira, T., Saitô, H., Katayama, M., Marumo, S., and Sugimura, T. 1-Nitroso-indole-3-acetonitrile, a mutagen produced by nitrite treatment of indole-3-acetonitrile. *Proc. Japan. Acad.* 61B: 190-192 (1985).
 73. Berenblum, I. The mechanisms of carcinogenesis. *Cancer Res.* 1: 807-814 (1941).
 74. Hecker, E. Phorbol ester from croton oil, chemical nature and biological activities. *Naturwiss.* 54: 282-284 (1967).
 75. Fujiki, H., Mori, M., Nakayasu, N., Terada, M., Sugimura, T., and Moore, R. E. Indole alkaloids: dihydroteleocidin B, teleocidin, and lyngbyatoxin A as members of a new class tumor promoters. *Proc. Natl. Acad. Sci. (U.S.)* 78: 3872-3876 (1981).
 76. Suganuma, M., Fujiki, H., Tahira, T., Cheuk, G., Moore, R. E., and Sugimura, T. Estimation of tumor promoting activity and structure-function relationships of aplysiatoxins. *Carcinogenesis* 5: 315-318 (1984).
 77. Boutwell, R. K. Some biological aspects of skin carcinogenesis. *Progr. Exptl. Tumor Res.* 4: 207-250 (1964).